

EXTREMAL GRAPH THEORY FOR METRIC DIMENSION AND DIAMETER

CARMEN HERNANDO, MERCÈ MORA, IGNACIO M. PELAYO, CARLOS SEARA,
AND DAVID R. WOOD

ABSTRACT. A set of vertices S *resolves* a connected graph G if every vertex is uniquely determined by its vector of distances to the vertices in S . The *metric dimension* of G is the minimum cardinality of a resolving set of G . Let $\mathcal{G}_{\beta,D}$ be the set of graphs with metric dimension β and diameter D . It is well-known that the minimum order of a graph in $\mathcal{G}_{\beta,D}$ is exactly $\beta + D$. The first contribution of this paper is to characterise the graphs in $\mathcal{G}_{\beta,D}$ with order $\beta + D$ for all values of β and D . Such a characterisation was previously only known for $D \leq 2$ or $\beta \leq 1$. The second contribution is to determine the maximum order of a graph in $\mathcal{G}_{\beta,D}$ for all values of D and β . Only a weak upper bound was previously known.

1. INTRODUCTION

Let G be a connected graph¹. A vertex $x \in V(G)$ *resolves*² a pair of vertices $v, w \in V(G)$ if $\text{dist}(v, x) \neq \text{dist}(w, x)$. A set of vertices $S \subseteq V(G)$ *resolves* G , and S is a *resolving set* of G , if every pair of distinct vertices of G are resolved by some vertex in S . Informally, S resolves G if every vertex of G is uniquely determined by its vector of distances to the

2000 *Mathematics Subject Classification.* 05C12 (distance in graphs), 05C35 (extremal graph theory).

Key words and phrases. graph, distance, resolving set, metric dimension, metric basis, diameter, order.

The research of Carmen Hernando, Mercè Mora, Carlos Seara, and David Wood is supported by the projects MEC MTM2006-01267 and DURSI 2005SGR00692. The research of Ignacio Pelayo is supported by the projects MTM2005-08990-C02-01 and SGR2005-00412. The research of David Wood is supported by a Marie Curie Fellowship of the European Community under contract 023865.

¹Graphs in this paper are finite, undirected, and simple. The vertex set and edge set of a graph G are denoted by $V(G)$ and $E(G)$. For vertices $v, w \in V(G)$, we write $v \sim w$ if $vw \in E(G)$, and $v \not\sim w$ if $vw \notin E(G)$. For $S \subseteq V(G)$, let $G[S]$ be the subgraph of G induced by S . That is, $V(G[S]) = S$ and $E(G[S]) = \{vw \in E(G) : v \in S, w \in S\}$. For $S \subseteq V(G)$, let $G \setminus S$ be the graph $G[V(G) \setminus S]$. For $v \in V(G)$, let $G \setminus v$ be the graph $G \setminus \{v\}$. Suppose that G is connected. The *distance* between vertices $v, w \in V(G)$, denoted by $\text{dist}_G(v, w)$, is the length (that is, the number of edges) in a shortest path between v and w in G . The *eccentricity* of a vertex v in G is $\text{ecc}_G(v) := \max\{\text{dist}_G(v, w) : w \in V(G)\}$. We drop the subscript G from these notations if the graph G is clear from the context. The *diameter* of G is $\text{diam}(G) := \max\{\text{dist}(v, w) : v, w \in V(G)\} = \max\{\text{ecc}(v) : v \in V(G)\}$. For integers $a \leq b$, let $[a, b] := \{a, a+1, \dots, b\}$.

²It will be convenient to also use the following definitions for a connected graph G . A vertex $x \in V(G)$ *resolves* a set of vertices $T \subseteq V(G)$ if x resolves every pair of distinct vertices in T . A set of vertices $S \subseteq V(G)$ *resolves* a set of vertices $T \subseteq V(G)$ if for every pair of distinct vertices $v, w \in T$, there exists a vertex $x \in S$ that resolves v, w .

vertices in S . A resolving set S of G with the minimum cardinality is a *metric basis* of G , and $|S|$ is the *metric dimension* of G , denoted by $\beta(G)$.

Resolving sets in general graphs were first defined by Slater [28] and Harary and Melter [15]. Resolving sets have since been widely investigated [2, 3, 4, 5, 6, 7, 9, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 29, 31, 32, 33], and arise in diverse areas including coin weighing problems [10, 14, 16, 18, 30], network discovery and verification [1], robot navigation [17, 27], connected joins in graphs [26], the Djoković-Winkler relation [3], and strategies for the Mastermind game [8, 11, 12, 13, 16].

For positive integers β and D , let $\mathcal{G}_{\beta,D}$ be the class of connected graphs with metric dimension β and diameter D . Consider the following two extremal questions:

- What is the minimum order of a graph in $\mathcal{G}_{\beta,D}$?
- What is the maximum order of a graph in $\mathcal{G}_{\beta,D}$?

The first question was independently answered by Yushmanov [33], Khuller et al. [17], and Chartrand et al. [5], who proved that the minimum order of a graph in $\mathcal{G}_{\beta,D}$ is $\beta + D$ (see Lemma 2.2). Thus it is natural to consider the following problem:

- Characterise the graphs in $\mathcal{G}_{\beta,D}$ with order $\beta + D$.

Such a characterisation is simple for $\beta = 1$. In particular, Khuller et al. [17] and Chartrand et al. [5] independently proved that paths P_n (with $n \geq 2$ vertices) are the only graphs with metric dimension 1. Thus $\mathcal{G}_{1,D} = \{P_{D+1}\}$.

The characterisation is again simple at the other extreme with $D = 1$. In particular, Chartrand et al. [5] proved that the complete graph K_n (with $n \geq 1$ vertices) is the only graph with metric dimension $n - 1$ (see Proposition 2.12). Thus $\mathcal{G}_{\beta,1} = \{K_{\beta+1}\}$.

Chartrand et al. [5] studied the case $D = 2$, and obtained a non-trivial characterisation of graphs in $\mathcal{G}_{\beta,2}$ with order $\beta + 2$ (see Proposition 2.13).

The first contribution of this paper is to characterise the graphs in $\mathcal{G}_{\beta,D}$ with order $\beta + D$ for all values of $\beta \geq 1$ and $D \geq 3$, thus completing the characterisation for all values of D . This result is stated and proved in Section 2.

We then study the second question above: What is the maximum order of a graph in $\mathcal{G}_{\beta,D}$? Previously, only a weak upper bound was known. In particular, Khuller et al. [17] and Chartrand et al. [5] independently proved that every graph in $\mathcal{G}_{\beta,D}$ has at most $D^\beta + \beta$ vertices. This bound is tight only for $D \leq 3$ or $\beta = 1$.

Our second contribution is to determine the (exact) maximum order of a graph in $\mathcal{G}_{\beta,D}$ for all values of D and β . This result is stated and proved in Section 3.

2. GRAPHS WITH MINIMUM ORDER

In this section we characterise the graphs in $\mathcal{G}_{\beta,D}$ with minimum order. We start with an elementary lemma.

Lemma 2.1. *Let S be a set of vertices in a connected graph G . Then $V(G) \setminus S$ resolves G if and only if every pair of vertices in S are resolved by some vertex not in S .*

Proof. If $v \in V(G) \setminus S$ and w is any other vertex, then v resolves v and w . By assumption every pair of vertices in S are resolved by some vertex in $V(G) \setminus S$. \square

Lemma 2.1 enables the minimum order of a graph in $\mathcal{G}_{\beta,D}$ to be easily determined.

Lemma 2.2 ([5, 17, 33]). *The minimum order of a graph in $\mathcal{G}_{\beta,D}$ is $\beta + D$.*

Proof. First we prove that every graph $G \in \mathcal{G}_{\beta,D}$ has order at least $\beta + D$. Let v_0, v_D be vertices such that $\text{dist}(v_0, v_D) = D$. Let $P = (v_0, v_1, \dots, v_D)$ be a path of length D in G . Then v_0 resolves v_i, v_j for all distinct $i, j \in [1, D]$. Thus $V(G) \setminus \{v_1, \dots, v_D\}$ resolves G by Lemma 2.1. Hence $\beta \leq |V(G)| - D$ and $|V(G)| \geq \beta + D$.

It remains to construct a graph $G \in \mathcal{G}_{\beta,D}$ with order $\beta + D$. Let G be the ‘broom’ tree obtained by adding β leaves adjacent to one endpoint of the path on D vertices. Observe that $|V(G)| = \beta + D$ and G has diameter D . It follows from Slater’s formula [28] for the metric dimension of a tree³ that the β leaves adjacent to one endpoint of the path are a metric basis of G . Hence $G \in \mathcal{G}_{\beta,D}$. \square

2.1. Twin Vertices. Let u be a vertex of a graph G . The *open neighborhood* of u is $N(u) := \{v \in V(G) : uv \in E(G)\}$, and the *closed neighborhood* of u is $N[u] := N(u) \cup \{u\}$. Two distinct vertices u, v are *adjacent twins* if $N[u] = N[v]$, and *non-adjacent twins* if $N(u) = N(v)$. Observe that if u, v are adjacent twins then $uv \in E(G)$, and if u, v are non-adjacent twins then $uv \notin E(G)$; thus the names are justified⁴. If u, v are adjacent or non-adjacent twins, then u, v are *twins*. The next lemma follows from the definitions.

Lemma 2.3. *If u, v are twins in a connected graph G , then $\text{dist}(u, x) = \text{dist}(v, x)$ for every vertex $x \in V(G) \setminus \{u, v\}$.* \square

Corollary 2.4. *Suppose that u, v are twins in a connected graph G and S resolves G . Then u or v is in S . Moreover, if $u \in S$ and $v \notin S$, then $(S \setminus \{u\}) \cup \{v\}$ also resolves G .* \square

Lemma 2.5. *In a set S of three vertices in a graph, it is not possible that two vertices in S are adjacent twins, and two vertices in S are non-adjacent twins.*

Proof. Suppose on the contrary that u, v are adjacent twins and v, w are non-adjacent twins. Since u, v are twins and $v \not\sim w$, we have $u \not\sim w$. Similarly, since v, w are twins and $u \sim v$, we have $u \sim w$. This is the desired contradiction. \square

Lemma 2.6. *Let u, v, w be distinct vertices in a graph. If u, v are twins and v, w are twins, then u, w are also twins.*

Proof. Suppose that u, v are adjacent twins. That is, $N[u] = N[v]$. By Lemma 2.5, v, w are adjacent twins. That is, $N[v] = N[w]$. Hence $N[u] = N[w]$. That is, u, w are adjacent twins. By a similar argument, if u, v are non-adjacent twins, then v, w are non-adjacent twins and u, w are non-adjacent twins. \square

³Also see [5, 15, 17] for proofs of Slater’s formula.

⁴In the literature, adjacent twins are called *true* twins, and non-adjacent twins are called *false* twins. We prefer the more descriptive names, *adjacent* and *non-adjacent*.

For a graph G , a set $T \subseteq V(G)$ is a *twin-set* of G if v, w are twins in G for every pair of distinct vertices $v, w \in T$.

Lemma 2.7. *If T is a twin-set of a graph G , then either every pair of vertices in T are adjacent twins, or every pair of vertices in T are non-adjacent twins.*

Proof. Suppose on the contrary some pair of vertices $v, w \in T$ are adjacent twins, and some pair of vertices $x, y \in T$ are adjacent twins. If v, x are adjacent twins then $\{v, x, y\}$ contradict Lemma 2.5. Otherwise v, x are non-adjacent twins, in which case $\{v, w, x\}$ contradict Lemma 2.5. \square

Lemma 2.8. *Let T be a twin-set of a connected graph G with $|T| \geq 3$. Then $\beta(G) = \beta(G \setminus u) + 1$ for every vertex $u \in T$.*

Proof. Let u, v, w be distinct vertices in T . By Corollary 2.4, there is a metric basis W of G such that $u, v \in W$. Since u has a twin in $G \setminus u$, for all $x, y \in V(G \setminus u)$ we have $\text{dist}_G(x, y) = \text{dist}_{G \setminus u}(x, y)$. In particular, $G \setminus u$ is connected. First we prove that $W \setminus \{u\}$ resolves $G \setminus u$. For all distinct vertices $x, y \in V(G \setminus u)$, there is a vertex $s \in W$ such that $\text{dist}_G(x, s) \neq \text{dist}_G(y, s)$. If $s \neq u$, then $s \in W \setminus \{u\}$ resolves the pair x, y . Otherwise, v is a twin of $s = u$ and $\text{dist}_{G \setminus u}(x, v) = \text{dist}_G(x, v) = \text{dist}_G(x, s) \neq \text{dist}_G(y, s) = \text{dist}_G(y, v) = \text{dist}_{G \setminus u}(y, v)$. Consequently, $v \in W \setminus \{u\}$ resolves the pair x, y . Now suppose that W' is a resolving set of $G \setminus u$ such that $|W'| < |W| - 1$. For all $x, y \in V(G \setminus u)$, there exists a vertex $s \in W'$ such that $\text{dist}_{G \setminus u}(x, s) \neq \text{dist}_{G \setminus u}(y, s)$. Then $W' \cup \{u\}$ is a resolving set in G of cardinality less than $|W|$, which contradicts the fact that W is a resolving set of minimum cardinality. \square

Note that it is necessary to assume that $|T| \geq 3$ in Lemma 2.8. For example, $\{x, z\}$ is a twin-set of the 3-vertex path $P_3 = (x, y, z)$, but $\beta(P_3) = \beta(P_3 \setminus x) = 1$.

Corollary 2.9. *Let T be a twin-set of a connected graph G with $|T| \geq 3$. Then $\beta(G) = \beta(G \setminus S) + |S|$ for every subset $S \subset T$ with $|S| \leq |T| - 2$.* \square

2.2. The Twin Graph. Let G be a graph. Define a relation \equiv on $V(G)$ by $u \equiv v$ if and only if $u = v$ or u, v are twins. By Lemma 2.6, \equiv is an equivalence relation. For each vertex $v \in V(G)$, let v^* be the set of vertices of G that are equivalent to v under \equiv . Let $\{v_1^*, \dots, v_k^*\}$ be the partition of $V(G)$ induced by \equiv , where each v_i is a representative of the set v_i^* . The *twin graph* of G , denoted by G^* , is the graph with vertex set $V(G^*) := \{v_1^*, \dots, v_k^*\}$, where $v_i^* v_j^* \in E(G^*)$ if and only if $v_i v_j \in E(G)$. The next lemma implies that this definition is independent of the choice of representatives.

Lemma 2.10. *Let G^* be the twin graph of a graph G . Then two vertices v^* and w^* of G^* are adjacent if and only if every vertex in v^* is adjacent to every vertex in w^* in G .*

Proof. Suppose on the contrary that some vertex in v^* is adjacent to some vertex in w^* , and some vertex in v^* is not adjacent to some vertex in w^* . Then $y \sim x \not\sim z$ for some vertices $x \in v^*$ and $y, z \in w^*$. Thus y, z are not twins, which is the desired contradiction. \square

Each vertex v^* of G^* is a maximal twin-set of G . By Lemma 2.7, $G[v^*]$ is a complete graph if the vertices of v^* are adjacent twins, or $G[v^*]$ is a null graph if the vertices of v^* are non-adjacent twins. So it makes sense to consider the following types of vertices in G^* . We say that $v^* \in V(G^*)$ is of *type*:

- (1) if $|v^*| = 1$,
 - (K) if $G[v^*] \cong K_r$ and $r \geq 2$,
 - (N) if $G[v^*] \cong N_r$ and $r \geq 2$,
- where N_r is the *null* graph with r vertices and no edges.

A vertex of G^* is of *type* (1K) if it is of type (1) or (K). A vertex of G^* is of *type* (1N) if it is of type (1) or (N). A vertex of G^* is of *type* (KN) if it is of type (K) or (N).

Observe that the graph G is uniquely determined by G^* , and the type and cardinality of each vertex of G^* . In particular, if v^* is adjacent to w^* in G^* , then every vertex in v^* is adjacent to every vertex in w^* in G .

We now show that the diameters of G and G^* are closely related.

Lemma 2.11. *Let $G \neq K_1$ be a connected graph. Then $\text{diam}(G^*) \leq \text{diam}(G)$. Moreover, $\text{diam}(G^*) < \text{diam}(G)$ if and only if $G^* \cong K_n$ for some $n \geq 1$. In particular, if $\text{diam}(G) \geq 3$ then $\text{diam}(G) = \text{diam}(G^*)$.*

Proof. If v, w are adjacent twins in G , then $\text{dist}_G(v, w) = 1$ and $v^* = w^*$. If v, w are non-adjacent twins in G , then (since G has no isolated vertices) $\text{dist}_G(v, w) = 2$ and $v^* = w^*$. If v, w are not twins, then there is a shortest path between v and w that contains no pair of twins (otherwise there is a shorter path); thus

$$(1) \quad \text{dist}_G(v, w) = \text{dist}_{G^*}(v^*, w^*).$$

This implies that $\text{diam}(G^*) \leq \text{diam}(G)$. Moreover, if $\text{ecc}_G(v) \geq 3$ then v is not a twin of every vertex w for which $\text{dist}_G(v, w) = \text{ecc}_G(v)$; thus $\text{dist}_G(v, w) = \text{dist}_{G^*}(v^*, w^*)$ by Equation (1) and $\text{ecc}_G(v) = \text{ecc}_{G^*}(v^*)$. Hence if $\text{diam}(G) \geq 3$ then $\text{diam}(G) = \text{diam}(G^*)$.

Now suppose that $\text{diam}(G) > \text{diam}(G^*)$. Thus $\text{diam}(G) \leq 2$. If $\text{diam}(G) = 1$ then G is a complete graph and $G^* \cong K_1$, as claimed. Otherwise $\text{diam}(G) = 2$ and $\text{diam}(G^*) \leq 1$; thus $G^* \cong K_n$ for some $n \geq 1$, as claimed.

It remains to prove that $\text{diam}(G^*) < \text{diam}(G)$ whenever $G^* \cong K_n$. In this case, $\text{diam}(G^*) \leq 1$. So we are done if $\text{diam}(G) \geq 2$. Otherwise $\text{diam}(G) \leq 1$ and G is also a complete graph. Thus $G^* \cong K_1$ and $\text{diam}(G^*) = 0$. Since $G \neq K_1$, we have $\text{diam}(G) = 1 > 0 = \text{diam}(G^*)$, as desired. \square

Note that graphs with $\text{diam}(G^*) < \text{diam}(G)$ include the complete multipartite graphs.

Theorem 2.14 below characterizes the graphs in $\mathcal{G}_{\beta, D}$ for $D \geq 3$ in terms of the twin graph. Chartrand et al. [5] characterized⁵ the graphs in $\mathcal{G}_{\beta, D}$ for $D \leq 2$. For consistency

⁵To be more precise, Chartrand et al. [5] characterised the graphs with $\beta(G) = n - 2$. By Lemma 2.2, if $\beta(G) = n - 2$ then G has diameter at most 2. By Proposition 2.12, if G has diameter 1 then $\beta(G) = n - 1$. Thus if $\beta(G) = n - 2$ then G has diameter 2.

with Theorem 2.14, we describe the characterisation by Chartrand et al. [5] in terms of the twin graph.

Proposition 2.12 ([5]). *The following are equivalent for a connected graph G with n vertices:*

- G has metric dimension $\beta(G) = n - 1$,
- $G \cong K_n$,
- $\text{diam}(G) = 1$,
- the twin graph G^* has one vertex, which is of type $(1K)$.

Proposition 2.13 ([5]). *The following are equivalent for a connected graph G with $n \geq 3$ vertices:*

- G has metric dimension $\beta(G) = n - 2$,
- G has metric dimension $\beta(G) = n - 2$ and diameter $\text{diam}(G) = 2$,
- the twin graph G^* of G satisfies
 - $G^* \cong P_2$ with at least one vertex of type (N) , or
 - $G^* \cong P_3$ with one leaf of type (1) , the other leaf of type $(1K)$, and the degree-2 vertex of type $(1K)$.

To describe our characterisation we introduce the following notation. Let $P_{D+1} = (u_0, u_1, \dots, u_D)$ be a path of length D . As illustrated in Figure 1(a), for $k \in [3, D - 1]$ let $P_{D+1,k}$ be the graph obtained from P_{D+1} by adding one vertex adjacent to u_{k-1} . As illustrated in Figure 1(b), for $k \in [2, D - 1]$ let $P'_{D+1,k}$ be the graph obtained from P_{D+1} by adding one vertex adjacent to u_{k-1} and u_k .




FIGURE 1. The graphs (a) $P_{D+1,k}$ and (b) $P'_{D+1,k}$.

Theorem 2.14. *Let G be a connected graph of order n and diameter $D \geq 3$. Let G^* be the twin graph of G . Let $\alpha(G^*)$ be the number of vertices of G^* of type (K) or (N) . Then $\beta(G) = n - D$ if and only if G^* is one of the following graphs:*

- (1) $G^* \cong P_{D+1}$ and one of the following cases hold (see Figure 2):
 - (a) $\alpha(G^*) \leq 1$;

- (b) $\alpha(G^*) = 2$, the two vertices of G^* not of type (1) are adjacent, and if one is a leaf of type (K) then the other is also of type (K);
 - (c) $\alpha(G^*) = 2$, the two vertices of G^* not of type (1) are at distance 2 and both are of type (N); or
 - (d) $\alpha(G^*) = 3$ and there is a vertex of type (N) or (K) adjacent to two vertices of type (N).
- (2) $G^* \cong P_{D+1,k}$ for some $k \in [3, D-1]$, the degree-3 vertex u_{k-1}^* of G^* is any type, each neighbour of u_{k-1}^* is type (1N), and every other vertex is type (1); see Figure 3.
- (3) $G^* \cong P'_{D+1,k}$ for some $k \in [2, D-1]$, the three vertices in the cycle are of type (1K), and every other vertex is of type (1); see Figure 4.




FIGURE 2. Cases (a)–(d) with $G^* \cong P_{D+1}$ in Theorem 2.14.





FIGURE 3. The case of $G^* \cong P_{D+1,k}$ in Theorem 2.14.


2.3. Proof of Necessity. Throughout this section, G is a graph of order n , diameter $D \geq 3$, and metric dimension $\beta(G) = n - D$. Let G^* be the twin graph of G .

Lemma 2.15. *There exists a vertex u_0 in G of eccentricity D with no twin.*

Proof. Let u_0 and u_D be vertices at distance D in G . As illustrated in Figure 5, let (u_0, u_1, \dots, u_D) be a shortest path between u_0 and u_D . Suppose on the contrary that both

FIGURE 4. The case of $G^* \cong P'_{D+1,k}$ in Theorem 2.14.

u_0 and u_D have twins. Let x be a twin of u_0 and y be a twin of u_D . We claim that $\{x, y\}$ resolves $\{u_0, \dots, u_D\}$. Now $u_0 \not\sim u_i$ for all $i \in [2, D]$, and thus $x \not\sim u_i$ (since x, u_0 are twins). Thus $\text{dist}(x, u_i) = i$ for each $i \in [1, D]$. Hence x resolves u_i, u_j for all distinct $i, j \in [1, D]$. By symmetry, $\text{dist}(y, u_i) = D - i$ for all $i \in [0, D - 1]$, and y resolves u_i, u_j for all distinct $i, j \in [0, D - 1]$. Thus $\{x, y\}$ resolves $\{u_0, \dots, u_D\}$, except for possibly the pair u_0, u_D . Now $\text{dist}(x, u_0) \leq 2$ and $\text{dist}(x, u_D) = D$. Since $D \geq 3$, x resolves u_0, u_D . Thus $\{x, y\}$ resolves $\{u_0, \dots, u_D\}$. By Lemma 2.1, $\beta(G) \leq n - (D + 1) < n - D$, which is a contradiction. Thus u_0 or u_D has no twin. \square

FIGURE 5. $\{x, y\}$ resolves $\{u_0, \dots, u_D\}$ in Lemma 2.15.

For the rest of the proof, fix a vertex u_0 of eccentricity D in G with no twin, which exists by Lemma 2.15. Thus $u_0^* = \{u_0\}$ and $\text{ecc}_{G^*}(u_0^*) = \text{ecc}_G(u_0) = D$, which is also the diameter of G^* by Lemma 2.11. As illustrated in Figure 6, for each $i \in [0, D]$, let


$$A_i^* := \{v^* \in V(G^*) : \text{dist}(u_0^*, v^*) = i\}, \text{ and}$$

$$A_i := \{v \in V(G) : \text{dist}(u_0, v) = i\} = \bigcup \{v^* : v^* \in A_i^*\}.$$


Note that the last equality is true because u_0 has no twin and $\text{dist}(u_0, v) = \text{dist}(u_0, w)$ if v, w are twins. For all $i \in [0, D]$, we have $|A_i| \geq 1$ and $|A_i^*| \geq 1$. Moreover, $|A_0| = |A_0^*| = 1$. Let (u_0, u_1, \dots, u_D) be a path in G such that $u_i \in A_i$ for each $i \in [0, D]$. Observe that if $v \in A_i$ is adjacent to $w \in A_j$ then $|i - j| \leq 1$. In particular, $(u_i, u_{i+1}, \dots, u_j)$ is a shortest path between u_i and u_j .

Lemma 2.16. *For each $k \in [1, D]$,*

- $G[A_k]$ is a complete graph or a null graph;
- $G^*[A_k^*]$ is a complete graph or a null graph, and all the vertices in A_k^* are of type $(1K)$ in the first case, and of type $(1N)$ in the second case.

FIGURE 6. The sets A_0, A_1, \dots, A_D .

Proof. Suppose that $G[A_k]$ is neither complete nor null for some $k \in [1, D]$. Thus there exist vertices $u, v, w \in A_k$ such that $u \sim v \not\sim w$, as illustrated in Figure 7⁶. Let $S := (\{u_1, \dots, u_D\} \setminus \{u_k\}) \cup \{u, w\}$. Every pair of vertices in S are resolved by u_0 , except for u and w which are resolved by v . Thus $\{u_0, v\}$ resolves S . By Lemma 2.1, $\beta(G) \leq n - (D + 1) < n - D$. This contradiction proves the first claim, which immediately implies the second claim. \square

FIGURE 7. $\{u_0, v\}$ resolves $(\{u_1, \dots, u_D\} \setminus \{u_k\}) \cup \{u, w\}$ in Lemma 2.16.

Lemma 2.17. *For each $k \in [1, D]$, if $|A_k| \geq 2$ then*

- (a) $v \sim w$ for all vertices $v \in A_{k-1}$ and $w \in A_k$;
- (b) $v^* \sim w^*$ for all vertices $v^* \in A_{k-1}^*$ and $w^* \in A_k^*$.

Proof. First we prove (a). Every vertex in A_1 is adjacent to u_0 , which is the only vertex in A_0 . Thus (a) is true for $k = 1$. Now assume that $k \geq 2$. Suppose on the contrary that $v \not\sim w$ for some $v \in A_{k-1}$ and $w \in A_k$. There exists a vertex $u \in A_{k-1}$ adjacent to w . As illustrated in Figure 8, if $w \neq u_k$ then $\{u_0, w\}$ resolves $(\{u_1, \dots, u_D\} \setminus \{u_{k-1}\}) \cup \{u, v\}$.

As illustrated in Figure 9, if $w = u_k$ then $v \neq u_{k-1}$ and there exists a vertex $z \neq u_k$ in A_k , implying $\{u_0, u_k\}$ resolves $(\{u_1, \dots, u_D\} \setminus \{u_k\}) \cup \{v, z\}$. In both cases, Lemma 2.1

⁶In Figures 7–22, a solid line connects adjacent vertices, a dashed line connects non-adjacent vertices, and a coil connects vertices that may or may not be adjacent.




FIGURE 8. In Lemma 2.17, $\{u_0, w\}$ resolves $(\{u_1, \dots, u_D\} \setminus \{u_{k-1}\}) \cup \{u, v\}$.

implies that $\beta(G) \leq n - D - 1$. This contradiction proves (a), which immediately implies (b). \square




FIGURE 9. In Lemma 2.17, $\{u_0, u_k\}$ resolves $(\{u_1, \dots, u_D\} \setminus \{u_k\}) \cup \{v, z\}$.

Lemma 2.18. *If $|A_i| \geq 2$ and $|A_j| \geq 2$ then $|i - j| \leq 2$. Thus there are at most three distinct subsets A_i, A_j, A_k each with cardinality at least 2.*

Proof. As illustrated in Figure 10, suppose on the contrary that $|A_i| \geq 2$ and $|A_j| \geq 2$ for some $i, j \in [1, D]$ with $j \geq i + 3$. Let $x \neq u_i$ be a vertex in A_i . Let $y \neq u_j$ be a vertex in A_j . We claim that $\{u_j, x\}$ resolves $(\{u_0, \dots, u_D\} \setminus \{u_j\}) \cup \{y\}$.

By Lemma 2.17, $u_{i-1} \sim x$ and $u_{j-1} \sim y$. Observe that $\text{dist}(u_j, y) \in \{1, 2\}$; $\text{dist}(u_j, u_{j-h}) = h$ for all $h \in [1, j]$; $\text{dist}(u_j, u_{j+h}) = h$ for all $h \in [1, D - j]$. Thus u_j resolves $(\{u_0, \dots, u_D\} \setminus \{u_j\}) \cup \{y\}$, except for the following pairs:

- u_{j-h}, u_{j+h} whenever $1 \leq h \leq j \leq D - h$;
- y, u_{j-1} and y, u_{j+1} if $\text{dist}(y, u_j) = 1$; and
- y, u_{j-2} and y, u_{j+2} if $\text{dist}(y, u_j) = 2$.

We claim that x resolves each of these pairs. By Lemma 2.17, there is a shortest path between x and u_{j-1} that passes through u_{j-2} . Let $r := \text{dist}(x, u_{j-2})$. Thus $\text{dist}(x, u_{j-1}) = r + 1$, $\text{dist}(x, y) = r + 2$, $\text{dist}(x, u_{j+1}) = r + 3$, and $\text{dist}(x, u_{j+2}) = r + 4$. Thus x resolves every pair of vertices in $\{u_{j-2}, u_{j-1}, y, u_{j+1}, u_{j+2}\}$. It remains to prove that x resolves u_{j-h}, u_{j+h} whenever $3 \leq h \leq j \leq D - h$. Observe that $\text{dist}(x, u_{j+h}) \geq j + h - i$. If $j - h \geq i$ then, since $(x, u_{i-1}, \dots, u_{j-h})$ is a path,

$$\text{dist}(x, u_{j-h}) \leq j - h - i + 2 < j + h - i \leq \text{dist}(x, u_{j+h}).$$

Otherwise $j - h \leq i - 1$, implying

$$\text{dist}(x, u_{j-h}) = i - (j - h) < j + h - i \leq \text{dist}(x, u_{j+h}).$$

In each case $\text{dist}(x, u_{j-h}) < \text{dist}(x, u_{j+h})$. Thus x resolves u_{j-h}, u_{j+h} .

Hence $\{u_j, x\}$ resolves $(\{u_0, \dots, u_D\} \setminus \{u_j\}) \cup \{y\}$. By Lemma 2.1, $\beta(G) \leq n - D - 1$ which is the desired contradiction. \square




FIGURE 10. $\{u_j, x\}$ resolves $(\{u_0, \dots, u_D\} \setminus \{u_j\}) \cup \{y\}$ in Lemma 2.18.

Lemma 2.19. $|A_1^*| = 1$ and $|A_D^*| = 1$.

Proof. Consider a vertex $v \in A_1$. Then $v \sim u_0$ and every other neighbour of v is in $A_1 \cup A_2$. By Lemma 2.16, $G[A_1]$ is complete or null. If every vertex in A_1 is adjacent to every vertex in A_2 , then A_1 is a twin-set, and $|A_1^*| = 1$ as desired.

Now assume that some vertex $v \in A_1$ is not adjacent to some vertex in A_2 . By Lemma 2.17, the only vertex in A_2 is u_2 , and $v \not\sim u_2$. If $G[A_1]$ is null then $\text{ecc}(v) > D$, and if $G[A_1]$ is complete then v and u_0 are twins. In both cases we have a contradiction.

If $|A_D| = 1$ then $|A_D^*| = 1$. Now assume that $|A_D| \geq 2$. The neighbourhood of every vertex in A_D is contained in $A_{D-1} \cup A_D$. By Lemma 2.17, every vertex in A_D is adjacent to every vertex in A_{D-1} . By Lemma 2.16, $G[A_D]$ is complete or null. Thus A_D is a twin-set, implying $|A_D^*| = 1$. \square

Lemma 2.20. For each $k \in [1, D - 1]$, distinct vertices $v, w \in A_k$ are twins if and only if they have the same neighbourhood in A_{k+1} .

Proof. The neighbourhood of both v and w is contained in $A_{k-1} \cup A_k \cup A_{k+1}$. By Lemma 2.17, both v and w are adjacent to every vertex in A_{k-1} . By Lemma 2.16, $G[A_k]$ is complete or null. Thus v and w are twins if and only if they have the same neighbourhood in A_{k+1} . \square

Lemma 2.21. For each $k \in [2, D]$,

- (a) if $|A_k| \geq 2$ then $|A_{k-1}^*| = 1$;
- (b) if $|A_k| = 1$ then $|A_{k-1}^*| \leq 2$.

Proof. Suppose that $|A_k| \geq 2$. If $|A_{k-1}| = 1$ then $|A_{k-1}^*| = 1$ as desired. Now assume that $|A_{k-1}| \geq 2$. Thus A_{k-1} is a twin-set by Lemma 2.20, implying $|A_{k-1}^*| = 1$. Now suppose that $|A_k| = 1$. If $|A_{k-1}| = 1$ then $|A_{k-1}^*| = 1$ and we are done. So assume that $|A_{k-1}| \geq 2$. By Lemma 2.20, the set of vertices in A_{k-1} that are adjacent to the unique vertex in A_k

is a maximal twin-set, and the set of vertices in A_{k-1} that are not adjacent to the unique vertex in A_k is a maximal twin-set (if it is not empty). Therefore $|A_{k-1}^*| \leq 2$. \square

Lemma 2.22. *For each $k \in [1, D]$, we have $|A_k^*| \leq 2$. Moreover, there are at most three values of k for which $|A_k^*| = 2$. Furthermore, if $|A_i^*| = 2$ and $|A_j^*| = 2$ then $|i - j| \leq 2$.*

Proof. Lemma 2.19 proves the result for $k = D$. Now assume that $k \in [1, D - 1]$. Suppose on the contrary that $|A_k^*| \geq 3$ for some $k \in [1, D]$. By the contrapositive of Lemma 2.21(a), $|A_{k+1}| = 1$. By Lemma 2.21(b), $|A_k^*| \leq 2$, which is the desired contradiction. The remaining claims follow immediately from Lemma 2.18. \square

Lemma 2.23. *Suppose that $|A_k^*| = 2$ for some $k \in [2, D - 1]$. Then $|A_{k+1}| = |A_{k+1}^*| = 1$, and exactly one of the two vertices of A_k^* is adjacent to the only vertex of A_{k+1}^* . Moreover, if $k \leq D - 2$ then $|A_{k+2}| = |A_{k+2}^*| = 1$.*

Proof. By the contrapositive of Lemma 2.21(a), $|A_{k+1}| = |A_{k+1}^*| = 1$. By Lemma 2.20, exactly one vertex in A_k^* is adjacent to the vertex in A_{k+1}^* . Now suppose that $k \leq D - 2$ but $|A_{k+2}| \geq 2$. As illustrated in Figure 11, let $x \neq u_{k+2}$ be a vertex in A_{k+2} . Let $y \neq u_k$ be a vertex in A_k , such that y, u_k are not twins, that is, $y \not\sim u_{k+1}$. By Lemma 2.17, $u_{k-1} \sim y$ and $u_{k+1} \sim x$. Thus $\{x, u_0\}$ resolves $\{u_1, \dots, u_D, y\}$. By Lemma 2.1, $\beta(G) \leq n - D - 1$, which is a contradiction. Hence $|A_{k+2}| = 1$, implying $|A_{k+2}^*| = 1$. \square




FIGURE 11. $\{x, u_0\}$ resolves $\{u_1, \dots, u_D, y\}$ in Lemma 2.23.

We now prove that the structure of the graph G^* is as claimed in Theorem 2.14.

Lemma 2.24. *Either $G^* \cong P_{D+1}$, $G^* \cong P_{D+1,k}$ for some $k \in [3, D - 1]$, or $G^* \cong P'_{D+1,k}$ for some $k \in [2, D - 1]$.*

Proof. By Lemma 2.22 each set A_k^* contains at most two vertices of G^* . Lemmas 2.19, 2.18 and 2.23 imply that $|A_k^*| = 2$ for at most one $k \in [0, D]$. If $|A_k^*| = 1$ for every $k \in [0, D]$ then $G^* \cong P_{D+1}$ as desired.

Now assume that $|A_k^*| = 2$ for exactly one $k \in [0, D]$. By Lemma 2.19, $k \in [2, D - 1]$. Let w^* be the vertex in A_k^* besides u_k^* . Then $w^* \sim u_{k-1}^*$ by Lemma 2.17. If $w^* \sim u_k^*$ then $G^* \cong P'_{D+1,k}$. Otherwise $w^* \not\sim u_k^*$. Then $G^* \cong P_{D+1,k}$. It remains to prove that in this case $k \neq 2$.

Suppose on the contrary that $G^* \cong P_{D+1,k}$ and $k = 2$. Thus $|A_2^*| = 2$. Say $A_2^* = \{u_2^*, w^*\}$, where $u_2^* \not\sim w^*$. By Lemma 2.23, $|A_3^*| = 1$. Thus $A_3^* = \{u_3^*\}$. Since $u_2^* \sim u_3^*$, by Lemma 2.20, $w^* \not\sim u_3^*$. Thus u_1^* is the only neighbour of w^* . Hence every vertex in w^* is a twin of u_0 , which contradicts the fact that u_0 has no twin. Thus $k \neq 2$ if $G^* \cong P_{D+1,k}$. \square

We now prove restrictions about the type of the vertices in G^* . To start with, Lemma 2.18 implies:

Corollary 2.25. *If $G^* \cong P_{D+1}$ then $\alpha(G^*) \leq 3$ and the distance between every pair of vertices not of type (1) is at most 2.*

Lemma 2.26. *Suppose that $G^* \cong P_{D+1}$ and $\alpha(G^*) = 2$. If the two vertices of G^* not of type (1) are adjacent, and one of them is a leaf of type (K), then the other is also of type (K).*

Proof. As illustrated in Figure 12, let x and y be twins of u_{D-1} and u_D respectively. By assumption $G[A_D]$ is a complete graph. Suppose on the contrary that $G[A_{D-1}]$ is a null graph. By Lemma 2.17, every vertex in A_D is adjacent to every vertex in A_{D-1} . Thus y resolves $\{u_0, \dots, u_D\}$, except for the pair u_{D-1}, u_D , which is resolved by x . Thus $\{x, y\}$ resolves $\{u_0, \dots, u_D\}$. By Lemma 2.1, $\beta(G) \leq n - D - 1$, which is a contradiction. Thus $G[A_{D-1}]$ is a complete graph. \square




FIGURE 12. $\{x, y\}$ resolves $\{u_0, \dots, u_D\}$ in Lemma 2.26.

Lemma 2.27. *Suppose that $G^* \cong P_{D+1}$ and for some $k \in [2, D-1]$, the vertices u_{k-1}^* and u_{k+1}^* of G^* are both not of type (1). Then u_{k-1}^* and u_{k+1}^* are both of type (N).*


Proof. Let x and y be twins of u_{k-1} and u_{k+1} respectively. Suppose on the contrary that one of u_{k-1}^* and u_{k+1}^* is of type (K). Without loss of generality u_{k-1}^* is of type (K), as illustrated in Figure 13. Thus $u_{k-1} \sim x$. We claim that $\{x, y\}$ resolves $\{u_0, u_1, \dots, u_D\}$.

Observe that x resolves every pair of vertices of $\{u_0, u_1, \dots, u_D\}$ except for:

- each pair of vertices in $\{u_{k-2}, u_{k-1}, u_k\}$, which are all resolved by y since $d(y, u_k) = 1$, $d(y, u_{k-1}) = 2$, and $d(y, u_{k-2}) = 3$; and
- the pairs $\{u_{k-j}, u_{k+j-2} : j \in [3, \min\{k, D+2-k\}]\}$, which are all resolved by y since $d(y, u_{k-j}) = j+1$, and

$$d(y, u_{k+j-2}) = \begin{cases} j-2 & \text{if } j \geq 4, \\ 1 \text{ or } 2 & \text{if } j = 3. \end{cases}$$

Hence $\{x, y\}$ resolves $\{u_0, u_1, \dots, u_D\}$. Thus Lemma 2.1 implies $\beta(G) \leq n - D - 1$, which is the desired contradiction. Hence u_{k-1}^* and u_{k+1}^* are both of type (N). \square

FIGURE 13. $\{x, y\}$ resolves $\{u_0, \dots, u_D\}$ in Lemma 2.27.

Corollary 2.25 and Lemmas 2.26 and 2.27 prove the necessity of the conditions in Theorem 2.14 when $G^* \cong P_{D+1}$.


Lemma 2.28. *Suppose that $G^* \cong P_{D+1,k}$ for some $k \in [3, D-1]$, where $A_k^* = \{u_k^*, w^*\}$ and $w^* \sim u_{k-1}^*$. Then u_{k-2}^* , u_k^* and w^* are type (1N), u_{k-1}^* is any type, and every other vertex is type (1).*

Proof. Since $u_k^* \not\sim w^*$, Lemma 2.16 implies that u_k^* and w^* are both type (1N). By Lemmas 2.18 and 2.23, the remaining vertices are of type (1) except, possibly u_{k-2}^* and u_{k-1}^* . Suppose that u_{k-2}^* is of type (K), as illustrated in Figure 14. Let x be a twin of u_{k-2} . Then $x \sim u_{k-1}$. We claim that $\{x, w\}$ resolves $\{u_0, u_1, \dots, u_D\}$.

Observe that x resolves every pair of vertices in $\{u_0, u_1, \dots, u_D\}$ except for:

- each pair of vertices in $\{u_{k-3}, u_{k-2}, u_{k-1}\}$, which are all resolved by w since $d(w, u_{k-1}) = 1$, $d(w, u_{k-2}) = 2$, and $d(w, u_{k-3}) = 3$; and
- the pairs $\{u_{k-2-j}, u_{k-2+j} : j \in [2, \min\{k-2, D-k+2\}]\}$, which are all resolved by w since $d(w, u_{k-2-j}) = j+2$ and $d(w, u_{k-2+j}) = j$.

Thus $\{x, w\}$ resolves $\{u_0, u_1, \dots, u_D\}$. Hence Lemma 2.1 implies that $\beta(G) \leq n - D - 1$, which is the desired contradiction. \square

FIGURE 14. $\{x, w\}$ resolves $\{u_0, \dots, u_D\}$ in Lemma 2.28.

Lemma 2.29. *Suppose that $G^* \cong P'_{D+1,k}$ for some $k \in [2, D-1]$, where $A_k^* = \{u_k^*, w^*\}$ and $u_{k-1}^* \sim w^* \sim u_k^*$. Then u_{k-1}^* , u_k^* and w^* are type (1K), and every other vertex is type (1).*

Proof. Since $u_k^* \sim w^*$, Lemma 2.16 implies that u_k^* and w^* are type (1K). By Lemmas 2.18 and 2.23, the remaining vertices are type (1) except possibly u_{k-2}^* and u_{k-1}^* .

Suppose on the contrary that u_{k-2}^* is type (K) or (N) , as illustrated in Figure 15. Let x be a twin of u_{k-2} . We claim that $\{x, w\}$ resolves $\{u_0, u_1, \dots, u_D\}$. Observe that w resolves every pair of vertices in $\{u_0, u_1, \dots, u_D\}$, except for pairs

$$\{u_{k-1-j}, u_{k+j} : j \in [0, \min\{k-1, D-k\}]\}.$$

These pairs are all resolved by x since $d(x, u_{k+j}) = j+2$ and

$$d(x, u_{k-1-j}) = \begin{cases} j-1 & \text{if } j \geq 2, \\ 1 \text{ or } 2 & \text{if } j = 1, \\ 1 & \text{if } j = 0. \end{cases}$$

Thus $\{x, w\}$ resolves $\{u_0, u_1, \dots, u_D\}$.




FIGURE 15. $\{x, w\}$ resolves $\{u_0, u_1, \dots, u_D\}$ in Lemma 2.29.

Suppose on the contrary that u_{k-1}^* is type (N) , as illustrated in Figure 16. Let y be a twin of u_{k-1} . We claim that $\{y, w\}$ resolves $\{u_0, u_1, \dots, u_D\}$. Observe that w resolves every pair of vertices in $\{u_0, u_1, \dots, u_D\}$, except for pairs

$$\{u_{k-1-j}, u_{k+j} : j \in [0, \min\{k-1, D-k\}]\}.$$

These pairs are all resolved by y since $d(y, u_{k+j}) = j+1$ and

$$d(y, u_{k-1-j}) = \begin{cases} j & \text{if } j \geq 1, \\ 2 & \text{if } j = 0. \end{cases}$$

Thus $\{y, w\}$ resolves $\{u_0, u_1, \dots, u_D\}$.




FIGURE 16. $\{y, w\}$ resolves $\{u_0, u_1, \dots, u_D\}$ in Lemma 2.29.

By Lemma 2.1, in each case $\beta(G) \leq n - D - 1$, which is the desired contradiction. \square

Observe that Lemmas 2.28 and 2.29 imply the necessity of the conditions in Theorem 2.14 when $G^* \cong P_{D+1,k}$ or $G^* \cong P'_{D+1,k}$. This completes the proof of the necessity of the conditions in Theorem 2.14.

2.4. Proof of Sufficiency. Let G be a graph with n vertices and $\text{diam}(G) \geq 3$. Let T be a twin-set of cardinality $r \geq 3$ in G . Let G' be the graph obtained from G by deleting all but two of the vertices in T . As in Lemma 2.11, $\text{diam}(G') = \text{diam}(G)$. Say G' has order n' . Then by Corollary 2.9, $\beta(G') = \beta(G) - (r - 2)$. Since $n' = n - (r - 2)$, we have that $\beta(G) = n - D$ if and only if $\beta(G') = n' - D$. Thus it suffices to prove the sufficiency in Theorem 2.14 for graphs G whose maximal twin-sets have at most two vertices. We assume in the remainder of this section that every twin-set in G has at most two vertices.

Suppose that the twin graph G^* of G is one of the graphs stated in Theorem 2.14. We need to prove that $\beta(G) = n - D$. Since $\beta(G) \leq n - D$ by Lemma 2.2, it suffices to prove that every subset of $n - D - 1$ vertices of G is not a resolving set. By Corollary 2.4, every resolving set contains at least one vertex in each twin-set of cardinality 2. Observe also that, since $\alpha(G^*)$ is the number of vertices of G^* not of type (1), we have that $\alpha(G^*) = n - |V(G^*)|$.

Case 1. $G^* \cong P_{D+1}$ with vertices $u_0^* \sim u_1^* \sim \dots \sim u_D^*$. We now prove that for each subcase stated in Theorem 2.14 every set of $n - D - 1 = n - |V(G^*)| = \alpha(G^*)$ vertices of G does not resolve G . Suppose on the contrary that W is a resolving set of G of cardinality $\alpha(G^*)$.

Case 1(a). $\alpha(G^*) \leq 1$: We need at least one vertex to resolve a graph G of order $n \geq 2$. So $\alpha(G^*) = 1$. Thus G is not a path, but Khuller et al. [17] and Chartrand et al. [5] independently proved that every graph with metric dimension 1 is a path, which is a contradiction.

Case 1(b)(i). $\alpha(G^*) = 2$, and u_k^*, u_{k+1}^* are not of type (1) for some $k \in [1, D - 2]$: As illustrated in Figure 17, consider vertices $x \neq u_k$ in u_k^* , and $y \neq u_{k+1}$ in u_{k+1}^* . By Corollary 2.4, we may assume that $W = \{x, y\}$.

Suppose that u_k^* is type (N). Then $x \not\sim u_k$, implying $\text{dist}(x, u_k) = \text{dist}(x, u_{k+2}) = 2$ and $\text{dist}(y, u_k) = \text{dist}(y, u_{k+2}) = 1$. Thus neither x nor y resolve u_k, u_{k+2} .

Suppose that u_{k+1}^* is type (N). Then $y \not\sim u_{k+1}$, implying $\text{dist}(x, u_{k-1}) = \text{dist}(x, u_{k+1}) = 1$ and $\text{dist}(y, u_{k-1}) = \text{dist}(y, u_{k+1}) = 2$. Thus neither x nor y resolve u_{k-1}, u_{k+1} .

Suppose that u_k^* and u_{k+1}^* are both type (K). Then $x \sim u_k$ and $y \sim u_{k+1}$, implying $\text{dist}(x, u_k) = \text{dist}(x, u_{k+1}) = 1$ and $\text{dist}(y, u_k) = \text{dist}(y, u_{k+1}) = 1$. Thus neither x nor y resolve u_k, u_{k+1} .

In each case we have a contradiction.

Case 1(b)(ii). $\alpha(G^*) = 2$, u_{D-1}^* is not type (1), and u_D^* is not type (1): As illustrated in Figure 18, consider $x \neq u_{D-1}$ in u_{D-1}^* and $y \neq u_D$ in u_D^* . By Corollary 2.4, we may assume that $W = \{x, y\}$.

First suppose that u_D^* is of type (N). Then $y \not\sim u_D$, implying $\text{dist}(x, u_{D-2}) = \text{dist}(x, u_D) = 1$ and $\text{dist}(y, u_{D-2}) = \text{dist}(y, u_D) = 2$. Thus neither x nor y resolve u_{D-2}, u_D , which is a contradiction.




FIGURE 17. In Case 1(b)(i).

Suppose that u_D^* and u_{D-1}^* are both type (K). Then $x \sim u_{D-1}$ and $y \sim u_D$, implying $\text{dist}(x, u_{D-1}) = \text{dist}(x, u_D) = 1$ and $\text{dist}(y, u_{D-1}) = \text{dist}(y, u_D) = 1$. Thus neither x nor y resolve u_{D-1}, u_D , which is a contradiction.






FIGURE 18. In Case 1(b)(ii).

Case 1(c). $\alpha(G^*) = 2$ and u_{k-1}^* is type (N), and u_{k+1}^* is type (N) for some $k \in [2, D-1]$: As illustrated in Figure 19, consider $x \neq u_{k-1}$ in u_{k-1}^* and $y \neq u_{k+1}$ in u_{k+1}^* . By Corollary 2.4, we may assume that $W = \{x, y\}$. Since $x \not\sim u_{k-1}$ and $y \not\sim u_{k+1}$, we have $\text{dist}(x, u_{k-1}) = \text{dist}(x, u_{k+1}) = 2$ and $\text{dist}(y, u_{k-1}) = \text{dist}(y, u_{k+1}) = 2$. Thus neither x nor y resolve u_{k-1}, u_{k+1} , which is a contradiction.


FIGURE 19. $\{x, y\}$ does not resolve u_{k-1}, u_{k+1} in Case 1(c).

Case 1(d). $\alpha(G^*) = 3$, u_{k-1}^* is type (N), u_k^* is type (K) or (N), and u_{k+1}^* is type (N) for some $k \in [2, D-1]$: As illustrated in Figure 20, consider $x \neq u_{k-1}$ in u_{k-1}^* , $y \neq u_k$ in u_k^* , and $z \neq u_{k+1}$ in u_{k+1}^* . By Corollary 2.4, we may assume that $W = \{x, y, z\}$. Now $x \not\sim u_{k-1}$ and $z \not\sim u_{k+1}$. Thus $\text{dist}(x, u_{k-1}) = \text{dist}(x, u_{k+1}) = 2$, $\text{dist}(y, u_{k-1}) = \text{dist}(y, u_{k+1}) = 1$, and $\text{dist}(z, u_{k-1}) = \text{dist}(z, u_{k+1}) = 2$. Thus $\{x, y, z\}$ does not resolve u_{k-1}, u_{k+1} , which is a contradiction.

Case 2. $G^* \cong P_{D+1,k}$ for some $k \in [3, D-1]$: Thus G^* is path $(u_0^*, u_1^*, \dots, u_D^*)$ plus one vertex w^* adjacent to u_{k-1}^* . As illustrated in Figure 21, suppose that every vertex of G^*

FIGURE 20. $\{x, y, z\}$ does not resolve u_{k-1}, u_{k+1} in Case 1(d).

is of type (1), except for u_{k-2}^* , u_k^* and w^* which are type (1N), and u_{k-1}^* which is of any type. In this case $n - D - 1 = \alpha(G^*) + 1$. Consequently, it suffices to prove that $\alpha(G^*) + 1$ vertices do not resolve G . Suppose there is a resolving set W in G of cardinality $\alpha(G^*) + 1$. By Corollary 2.4, we can assume that W contains the $\alpha(G^*)$ twins of u_{k-2} , u_{k-1} , u_k and w (if they exist), and another vertex of G . Let x_{k-2} , x_{k-1} , x_k and y respectively be twin vertices of u_{k-2} , u_{k-1} , u_k and w (if they exist). Then $x_{k-2} \not\sim u_{k-2}$, $x_k \not\sim u_k$, and $y \not\sim w$. Thus the distance from x_{k-2} (respectively x_{k-1} , x_k , y) to any vertex of u_{k-2} , u_k , w is 2 (respectively 1, 2, 2). Hence any set of twins of vertices in $\{u_{k-2}, u_k, w\}$ (if they exist) does not resolve $\{u_{k-2}, u_k, w\}$. Moreover, if $i \in [0, k-1]$ then u_i does not resolve u_k, w ; if $i \in [k-1, D]$ then u_i does not resolve u_{k-2}, w ; and w does not resolve u_{k-2}, u_k . Therefore, $\alpha(G^*) + 1$ vertices do not resolve G .

FIGURE 21. $\{x_{k-2}, x_{k-1}, x_k, y\}$ does not resolve $\{u_{k-2}, u_k, w\}$ in Case 2.

Case 3. $G^* \cong P'_{D+1,k}$ for some $k \in [2, D-1]$: Thus G^* is path $(u_0^*, u_1^*, \dots, u_D^*)$ plus one vertex w^* adjacent to u_{k-1}^* and u_k^* . As illustrated in Figure 22, suppose that every vertex of G^* is type (1) except for u_{k-1}^* , u_k^* , and w^* which are of type (1K). In this case, $n - D - 1 = \alpha(G^*) + 1$. Consequently, it suffices to prove that $\alpha(G^*) + 1$ vertices do not resolve G . Suppose there is a resolving set W in G of cardinality $\alpha(G^*) + 1$. By Corollary 2.4, we may assume that W contains exactly the $\alpha(G^*)$ twin vertices of u_{k-1} , u_k and w (if they exist), and another vertex of G . Let x_{k-1} , x_k , and y respectively be twins of u_{k-1} , u_k and w (if they exist). Hence $x_{k-1} \sim u_{k-1}$, $x_k \sim u_k$ and $y \sim w$. Consequently, u_{k-1} , u_k and w are at distance 1 from x_k , x_{k+1} and y . Thus any set of twins of vertices in

$\{u_{k-1}, u_k, w\}$ (if they exist) does not resolve $\{u_{k-1}, u_k, w\}$. Moreover, if $i \in [0, k-1]$ then u_i does not resolve u_k, w ; if $i \in [k, D]$ then u_i does not resolve u_{k-1}, w ; and w does not resolve u_{k-1}, u_k . Thus $\alpha(G^*) + 1$ vertices do not resolve G . \square




FIGURE 22. $\{x_{k-1}, x_k, y\}$ does not resolve $\{u_{k-1}, u_k, w\}$ in Case 3.

3. GRAPHS WITH MAXIMUM ORDER

In this section we determine the maximum order of a graph in $\mathcal{G}_{\beta, D}$.

Theorem 3.1. *For all integers $D \geq 2$ and $\beta \geq 1$, the maximum order of a connected graph with diameter D and metric dimension β is*

$$(2) \quad \left(\left\lfloor \frac{2D}{3} \right\rfloor + 1 \right)^\beta + \beta \sum_{i=1}^{\lceil D/3 \rceil} (2i-1)^{\beta-1}.$$

First we prove the upper bound in Theorem 3.1.

Lemma 3.2. *For every graph $G \in \mathcal{G}_{\beta, D}$,*

$$|V(G)| \leq \left(\left\lfloor \frac{2D}{3} \right\rfloor + 1 \right)^\beta + \beta \sum_{i=1}^{\lceil D/3 \rceil} (2i-1)^{\beta-1}.$$

Proof. Let S be a metric basis of G . Let $k \in [0, D]$ be specified later. For each vertex $v \in S$ and integer $i \in [0, k]$, let $N_i(v) := \{x \in V(G) : \text{dist}(v, x) = i\}$.

Consider two vertices $x, y \in N_i(v)$. There is a path from x to v of length i , and there is a path from y to v of length i . Thus $\text{dist}(x, y) \leq 2i$. Hence for each vertex $u \in S$, the difference between $\text{dist}(u, x)$ and $\text{dist}(u, y)$ is at most $2i$. Thus the distance vector of x with respect to S has an i in the coordinate corresponding to v , and in each other coordinate, there are at most $2i+1$ possible values. Therefore $|N_i(v)| \leq (2i+1)^{\beta-1}$.

Consider a vertex $x \in V(G)$ that is not in $N_i(v)$ for all $v \in S$ and $i \in [0, k]$. Then $\text{dist}(x, v) \geq k+1$ for all $v \in S$. Thus the distance vector of x with respect to S consists of

β numbers in $[k+1, D]$. Thus there are at most $(D-k)^\beta$ such vertices. Hence

$$\begin{aligned} |V(G)| &\leq (D-k)^\beta + \sum_{v \in S} \sum_{i=0}^k |N_i(v)| \\ &\leq (D-k)^\beta + \beta \sum_{i=0}^k (2i+1)^{\beta-1} . \end{aligned}$$

Note that with $k=0$ we obtain the bound $|V(G)| \leq D^\beta + \beta$, independently due to Khuller et al. [17] and Chartrand et al. [5]. Instead we define $k := \lceil D/3 \rceil - 1$. Then $k \in [0, D]$ and

$$\begin{aligned} |V(G)| &\leq \left(D - \left\lfloor \frac{D}{3} \right\rfloor + 1 \right)^\beta + \beta \sum_{i=0}^{\lceil D/3 \rceil - 1} (2i+1)^{\beta-1} \\ &= \left(\left\lfloor \frac{2D}{3} \right\rfloor + 1 \right)^\beta + \beta \sum_{i=1}^{\lceil D/3 \rceil} (2i-1)^{\beta-1} . \end{aligned}$$

□

To prove the lower bound in Theorem 3.1 we construct a graph $G \in \mathcal{G}_{\beta,D}$ with as many vertices as in Equation (2). The following definitions apply for the remainder of this section. Let $A := \lceil D/3 \rceil$ and $B := \lceil D/3 \rceil + \lfloor D/3 \rfloor$. Consider the following subsets of \mathbb{Z}^β . Let

$$Q := \{(x_1, \dots, x_\beta) : A \leq x_i \leq D, i \in [1, \beta]\} .$$

For each $i \in [1, \beta]$ and $r \in [0, A-1]$, let

$$P_{i,r} := \{(x_1, \dots, x_{i-1}, r, x_{i+1}, \dots, x_\beta) : x_j \in [B-r, B+r], j \neq i\} .$$

Let $P_i := \bigcup \{P_{i,r} : r \in [0, A-1]\}$ and $P := \bigcup \{P_i : i \in [1, \beta]\}$. Let G be the graph with vertex set $V(G) := Q \cup P$, where two vertices (x_1, \dots, x_β) and (y_1, \dots, y_β) in $V(G)$ are adjacent if and only if $|y_i - x_i| \leq 1$ for each $i \in [1, \beta]$. Figures 23 and 24 illustrate G for $\beta = 2$ and $\beta = 3$ respectively.





FIGURE 23. The graph G with $\beta = 2$. The shaded regions are Q , P_1 , and P_2 .

FIGURE 24. The convex hull of $V(G)$ with $\beta = 3$.

Lemma 3.3. *For all positive integers D and β ,*

$$|V(G)| = \left(\left\lfloor \frac{2D}{3} \right\rfloor + 1 \right)^\beta + \beta \sum_{i=1}^{\lceil D/3 \rceil} (2i-1)^{\beta-1}.$$

Proof. Observe that each coordinate of each vertex in Q is at least A , and each vertex in P has some coordinate less than A . Thus $Q \cap P = \emptyset$. Each vertex in P_j ($j \neq i$) has an i -coordinate at least $B - r \geq B - (A - 1) = \lfloor D/3 \rfloor + 1$, and each vertex in P_i has an i -coordinate of $r \leq A - 1 < \lfloor D/3 \rfloor + 1$. Thus $P_i \cap P_j = \emptyset$ whenever $i \neq j$. Each vertex in $P_{i,r}$ has an i -coordinate of r . Thus $P_{i,r} \cap P_{i,s} = \emptyset$ whenever $r \neq s$. Thus

$$\begin{aligned} |V(G)| &= |Q| + \sum_{i=1}^{\beta} \sum_{r=0}^{A-1} |P_{i,r}| \\ &= \left(D - \left(\left\lfloor \frac{D}{3} \right\rfloor - 1 \right) \right)^\beta + \beta \sum_{r=0}^{A-1} (2r+1)^{\beta-1} \\ &= \left(\left\lfloor \frac{2D}{3} \right\rfloor + 1 \right)^\beta + \beta \sum_{r=1}^A (2r-1)^{\beta-1}. \end{aligned}$$

□

We now determine the diameter of G . For distinct vertices $x = (x_1, \dots, x_\beta)$ and $y = (y_1, \dots, y_\beta)$ of G , let $z(x, y) := (z_1, \dots, z_\beta)$ where

$$z_i = \begin{cases} x_i & \text{if } x_i = y_i, \\ x_i + 1 & \text{if } x_i < y_i, \\ x_i - 1 & \text{if } x_i > y_i. \end{cases}$$

Lemma 3.4. $z(x, y) \in V(G)$ for all distinct vertices $x, y \in V(G)$.

Proof. The following observations are an immediate consequence of the definition of $z(x, y)$, where $h, k \in \mathbb{Z}$ and $j \in [1, \beta]$:

- (i) if $x_j, y_j \in [h, k]$ then $z_j \in [h, k]$;

- (ii) if $x_j \in [h, k]$ then $z_j \in [h - 1, k + 1]$; and
- (iii) if $x_j \in [h, k]$ and $y_j \in [h', k']$ for some $h' > h$ and $k' < k$, then $z_j \in [h + 1, k - 1]$.

We distinguish the following cases:

- (1) $x, y \in Q$: Then $x_j, y_j \in [A, D]$ for all j . Thus $z_j \in [a, D]$ by (i). Hence $z \in Q$.
- (2) $x \in P$ and $y \in Q$: Without loss of generality, $x \in P_{1,r}$; that is, $x = (r, x_2, \dots, x_\beta)$, where $r \in [0, A - 1]$ and $x_j \in [B - r, B + r]$. Since $y \in Q$, we have $y_1 \geq A > r$. Thus $z = (r + 1, z_2, \dots, z_\beta)$.
 - (2.1) $z_1 = r + 1 < A$: By (ii), $z_j \in [B - r - 1, B + r + 1]$ for every $j \neq 1$. Thus $z \in P_{1,r+1}$.
 - (2.2) $z_1 = r + 1 = A$: Then $z_1 \in [A, D]$. On the other hand, if $j \neq 2$ then $y_j \geq A$, and since $x \in P_{1,r}$, we have $r = A - 1$, we have $x_j \geq B - r = \lfloor D/3 \rfloor + 1 \geq \lceil D/3 \rceil = A$. Thus $z_j \geq A$. Hence $z \in Q$.
- (3) $x \in Q$ and $y \in P$: Without loss of generality, $y \in P_{1,r}$. That is, $y = (r, y_2, \dots, y_\beta)$, where $r \in [0, A - 1]$, $y_j \in [B - r, B + r]$, and $x = (x_1, \dots, x_\beta)$, where $x_j \geq A$ for all j . That is, $x_1 > r = y_1$, and therefore $z = (x_1 - 1, z_2, \dots, z_\beta)$.
 - (3.1) $z_1 = x_1 - 1 \geq A$: Since $x_j, y_j \geq A$, by (i), $z_j \geq A$ for every $j \neq 1$. That is, $z \in Q$.
 - (3.2) $z_1 = x_1 - 1 = A - 1$: Since $r \leq A - 1$, we have $y_j \in [B - r, B + r] \subseteq [B - (A - 1), B + A - 1]$ for all $j \notin \{1, 2\}$. Now $B - A = \lfloor D/3 \rfloor \leq \lceil D/3 \rceil = A$ and $D \leq \lfloor D/3 \rfloor + 2\lceil D/3 \rceil = A + B$. Thus $x_j \in [B - A, B + A]$. By (iii), $z_j \in [B - (A - 1), B + A - 1]$. That is, $z \in P_{1,A-1}$.
- (4) $x, y \in P_h$: Without loss of generality, $x, y \in P_1$. Thus $x = (r, x_2, \dots, x_\beta)$ for some $r \in [0, A - 1]$ with $x_j \in [B - r, B + r]$ for all $j \neq 1$, and $y = (s, y_2, \dots, y_\beta)$ for some $s \in [0, A - 1]$ with $y_j \in [B - s, B + s]$ for all $j \neq 1$.
 - (4.1) $r = s$: Then $z = (r, z_2, \dots, z_\beta)$. By (i), $z_j \in [B - r, B + r]$ for all $j \neq 1$. Thus $z \in P_{1,r}$.
 - (4.2) $r < s$: Then $z = (r + 1, z_2, \dots, z_\beta)$. By (ii), $z_j \in [B - (r + 1), B + r + 1]$ for all $j \neq 1$. Thus $z \in P_{1,r+1}$.
 - (4.3) $r > s$: Then $z = (r - 1, z_2, \dots, z_\beta)$. By (iii), $z_j \in [B - (r - 1), B + r - 1]$. Thus $z \in P_{1,r-1}$.
- (5) $x \in P_h, y \in P_k$ and $h \neq k$: Without loss of generality, $x \in P_1$ and $y \in P_2$. Thus $x = (r, x_2, \dots, x_\beta)$ for some $r \in [0, A - 1]$ with $x_j \in [B - r, B + r]$ for all $j \neq 1$, and $y = (y_1, s, y_3, \dots, y_\beta)$ for some $s \in [0, A - 1]$ with $y_j \in [B - s, B + s]$ for all $j \neq 2$. Hence $r < A \leq y_1$ and $s < A \leq x_2$, implying $z = (r + 1, x_2 - 1, z_3, \dots, z_\beta)$.
 - (5.1) $z_1 = r + 1 < A$: Now $x_j \in [B - r, B + r]$ for $j \neq 1$. Thus $z_j \in [B - r - 1, B + r + 1]$ by (ii). Thus $z \in P_{1,r+1}$.
 - (5.2) $z_1 = r + 1 = A$: Consider the following subcases:
 - (5.2.1) $z_2 = x_2 - 1 \geq A$: By hypotheses, $z_1, z_2 \geq A$. For $j \notin \{1, 2\}$, since $x_j, y_j \geq A$, (i) implies that $z_j \geq A$. Thus $z \in Q$.
 - (5.2.2) $z_2 = x_2 - 1 = A - 1$: In this case $x = (A - 1, A, x_3, \dots, x_\beta)$, $z = (A, A - 1, z_3, \dots, z_\beta)$, and $s \leq A - 1 = r$. Since $x \in P_{1,A-1}$, we have

$z_1 = x_2 = A \in [B - (A - 1), B + A - 1]$. For $j \notin \{1, 2\}$, since $x_j \in [B - (A - 1), B + A - 1]$ and $y_j \in [B - s, B + s]$, where $s \leq r = A - 1$, (i) implies that $z_j \in [B - (A - 1), B + A - 1]$. That is, $z \in P_{2, A-1}$. \square

Lemma 3.5. *For all vertices $x = (x_1, \dots, x_\beta)$ and $y = (y_1, \dots, y_\beta)$ of G ,*

$$\text{dist}(x, y) = \max\{|y_i - x_i| : i \in [1, \beta]\} \leq D.$$

Proof. For each $i \in [1, \beta]$, $\text{dist}(x, y) \geq |x_i - y_i|$ since on every xy -path P , the i -coordinates of each pair of adjacent vertices in P differ by at most 1. This proves the lower bound $\text{dist}(x, y) \geq \max_i |y_i - x_i|$.

Now we prove the upper bound $\text{dist}(x, y) \leq \max_i |y_i - x_i|$ by induction. If $\max_i |y_i - x_i| = 1$ then x and y are adjacent, and thus $\text{dist}(x, y) = 1$. Otherwise, let $z := z(x, y)$. By the definition of $z(x, y)$, for all $i \in [1, \beta]$ we have $|y_i - z_i| = |y_i - x_i| - 1$ unless $x_i = y_i$. Thus $\max_i |y_i - z_i| = \max_i |y_i - x_i| - 1$. By induction, $\text{dist}(z, y) \leq \max_i |y_i - z_i| = \max_i |y_i - x_i| - 1$. By Lemma 3.4, z is a vertex of G , and by construction, x and z are adjacent. Thus $\text{dist}(x, y) \leq \text{dist}(z, y) + 1 = \max_i |y_i - x_i|$, as desired. \square

Lemma 3.5 implies that G has diameter D . Let $S := \{v_1, \dots, v_\beta\}$, where

$$v_i = (\underbrace{t, \dots, t}_{i-1}, 0, t, \dots, t) .$$

Observe that each $v_i \in P_i$. We now prove that S is a metric basis of G .

Lemma 3.6. *$\text{dist}(x, v_i) = x_i$ for every vertex $x = (x_1, \dots, x_\beta)$ of G and for each $v_i \in S$.*

Proof. Let $v_{i,j}$ be the j -th coordinate of v_i ; that is, $v_{i,i} = 0$ and $v_{i,j} = t$ for $i \neq j$. Then $\text{dist}(x, v_i) = \max\{|v_{i,j} - x_j| : 1 \leq j \leq \beta\} = \max\{x_i, \max\{|t - x_j| : 1 \leq j \leq \beta, j \neq i\}\}$. We claim that $|t - x_j| \leq x_i$ for each $j \neq i$, implying $\text{dist}(x, v_i) = x_i$, as desired.

First suppose that $x \in Q$. Then $s \leq x_j \leq D$. Thus $|B - x_j| \leq \max\{B - A, D - t\} = \max\{\lfloor D/3 \rfloor, D - t\} \leq \max\{\lfloor D/3 \rfloor, \lceil D/3 \rceil\} = \lceil D/3 \rceil \leq x_i$.

Now suppose that $x \in P_{k,r}$ for some $k \neq i$ and for some r . Then $x_i \geq t - r \geq t - (\lceil D/3 \rceil - 1) = \lfloor D/3 \rfloor + 1 \geq \lceil D/3 \rceil$. Now $|t - x_j| \leq r \leq \lceil D/3 \rceil - 1$. Thus $|t - x_j| \leq x_i$.

Finally suppose that $x \in P_{i,r}$ for some r . Then $|t - x_j| \leq r = x_i$. \square

Lemma 3.6 implies that the metric coordinates of a vertex $x \in V(G)$ with respect to S are its coordinates as elements of \mathbb{Z}^β . Therefore S resolves G . Thus G has metric dimension at most $|S| = \beta$.

If the metric dimension of G was less than β , then by Lemma 3.2,

$$\left(\left\lfloor \frac{2D}{3} \right\rfloor + 1 \right)^\beta + \beta \sum_{i=1}^{\lceil D/3 \rceil} (2i-1)^{\beta-1} = |V(G)| \leq \left(\left\lfloor \frac{2D}{3} \right\rfloor + 1 \right)^{\beta-1} + (\beta-1) \sum_{i=1}^{\lceil D/3 \rceil} (2i-1)^{\beta-2} ,$$

which is a contradiction. Thus G has metric dimension β , and $G \in \mathcal{G}_{\beta,D}$. This completes the proof of Theorem 3.1.

REFERENCES

- [1] ZUZANA BEERLIOVA, FELIX EBERHARD, THOMAS ERLEBACH, ALEXANDER HALL, MICHAEL HOFFMANN, MATÚŠ MIHAL'ÁK, AND L. SHANKAR RAM. Network discovery and verification. *IEEE J. on Selected Areas in Communications*, 24(12):2168–2181, 2006.
- [2] ROBERT C. BRIGHAM, GARY CHARTRAND, RONALD D. DUTTON, AND PING ZHANG. Resolving domination in graphs. *Math. Bohem.*, 128(1):25–36, 2003.
- [3] JOSÉ CÁCERES, CARMEN HERNANDO, MERCÈ MORA, IGNACIO M. PELAYO, MARÍA L. PUERTAS, CARLOS SEARA, AND DAVID R. WOOD. On the metric dimension of cartesian products of graphs. *SIAM J. Discrete Math.*, to appear. arXiv.org/math/0507527.
- [4] GLENN G. CHAPPELL, JOHN GIMBEL, AND CHRIS HARTMAN. Bounds on the metric and partition dimensions of a graph, 2003. <http://www.cs.uaf.edu/~chappell/papers/metric/>.
- [5] GARY CHARTRAND, LINDA EROH, MARK A. JOHNSON, AND ORTRUD R. OELLERMANN. Resolvability in graphs and the metric dimension of a graph. *Discrete Appl. Math.*, 105(1-3):99–113, 2000.
- [6] GARY CHARTRAND, CHRISTOPHER POISSON, AND PING ZHANG. Resolvability and the upper dimension of graphs. *Comput. Math. Appl.*, 39(12):19–28, 2000.
- [7] GARY CHARTRAND AND PING ZHANG. The theory and applications of resolvability in graphs. A survey. In *Proc. 34th Southeastern International Conf. on Combinatorics, Graph Theory and Computing*, vol. 160 of *Congr. Numer.*, pp. 47–68. 2003.
- [8] VAŠEK CHVÁTAL. Mastermind. *Combinatorica*, 3(3-4):325–329, 1983.
- [9] JAMES CURRIE AND ORTRUD R. OELLERMANN. The metric dimension and metric independence of a graph. *J. Combin. Math. Combin. Comput.*, 39:157–167, 2001.
- [10] PAUL ERDŐS AND ALFRÉD RÉNYI. On two problems of information theory. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 8:229–243, 1963.
- [11] WAYNE GODDARD. Static mastermind. *J. Combin. Math. Combin. Comput.*, 47:225–236, 2003.
- [12] WAYNE GODDARD. Mastermind revisited. *J. Combin. Math. Combin. Comput.*, 51:215–220, 2004.
- [13] DON L. GREENWELL. Mastermind. *J. Recr. Math.*, 30:191–192, 1999-2000.
- [14] RICHARD K. GUY AND RICHARD J. NOWAKOWSKI. Coin-weighing problems. *Amer. Math. Monthly*, 102(2):164, 1995.
- [15] FRANK HARARY AND ROBERT A. MELTER. On the metric dimension of a graph. *Ars Combinatoria*, 2:191–195, 1976.
- [16] GRIGORI KABATIANSKI, V. S. LEBEDEV, AND J. THORPE. The Mastermind game and the rigidity of Hamming spaces. In *Proc. IEEE International Symposium on Information Theory (ISIT '00)*, p. 375. IEEE, 2000.

- [17] SAMIR KHULLER, BALAJI RAGHAVACHARI, AND AZRIEL ROSENFELD. Landmarks in graphs. *Discrete Appl. Math.*, 70(3):217–229, 1996.
- [18] BERNT LINDSTRÖM. On a combinatorial detection problem. I. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 9:195–207, 1964.
- [19] PAUL MANUEL, BHARATI RAJAN, INDRA RAJASINGH, AND M. CHRIS MONICA. Landmarks in torus networks. *J. Discrete Math. Sci. Cryptogr.*, 9(2):263–271, 2006.
- [20] JOEL PETERS-FRANSEN AND ORTRUD R. OELLERMANN. The metric dimension of the cartesian product of graphs. *Utilitas Math.*, 69:33–41, 2006.
- [21] CHRISTOPHER POISSON AND PING ZHANG. The metric dimension of unicyclic graphs. *J. Combin. Math. Combin. Comput.*, 40:17–32, 2002.
- [22] VARAPORN SAENPHOLPHAT AND PING ZHANG. Connected resolvability of graphs. *Czechoslovak Math. J.*, 53(128)(4):827–840, 2003.
- [23] VARAPORN SAENPHOLPHAT AND PING ZHANG. Conditional resolvability in graphs: a survey. *Int. J. Math. Math. Sci.*, (37-40):1997–2017, 2004.
- [24] VARAPORN SAENPHOLPHAT AND PING ZHANG. Detour resolvability of graphs. In *Proc. of 35th Southeastern International Conf. on Combinatorics, Graph Theory and Computing*, vol. 169 of *Congr. Numer.*, pp. 3–21. Utilitas Math., 2004.
- [25] VARAPORN SAENPHOLPHAT AND PING ZHANG. On connected resolving decompositions in graphs. *Czechoslovak Math. J.*, 54(129)(3):681–696, 2004.
- [26] ANDRÁS SEBŐ AND ERIC TANNIER. On metric generators of graphs. *Math. Oper. Res.*, 29(2):383–393, 2004.
- [27] B. SHANMUKHA, B. SOORYANARAYANA, AND K. S. HARINATH. Metric dimension of wheels. *Far East J. Appl. Math.*, 8(3):217–229, 2002.
- [28] PETER J. SLATER. Leaves of trees. In *Proc. 6th Southeastern Conf. on Combinatorics, Graph Theory, and Computing*, vol. 14 of *Congr. Numer.*, pp. 549–559. Utilitas Math., 1975.
- [29] PETER J. SLATER. Dominating and reference sets in a graph. *J. Math. Phys. Sci.*, 22(4):445–455, 1988.
- [30] STAFFAN SÖDERBERG AND HAROLD S. SHAPIRO. A combinatorial detection problem. *Amer. Math. Monthly*, 70:1066, 1963.
- [31] B. SOORYANARAYANA. On the metric dimension of a graph. *Indian J. Pure Appl. Math.*, 29(4):413–415, 1998.
- [32] B. SOORYANARAYANA AND B. SHANMUKHA. A note on metric dimension. *Far East J. Appl. Math.*, 5(3):331–339, 2001.
- [33] S. V. YUSHMANOV. Estimates for the metric dimension of a graph in terms of the diameters and the number of vertices. *Vestnik Moskov. Univ. Ser. I Mat. Mekh.*, 103:68–70, 1987.

DEPARTAMENT DE MATEMÀTICA APLICADA I, UNIVERSITAT POLITÈCNICA DE CATALUNYA, BARCELONA,
SPAIN

E-mail address: `carmen.hernando@upc.edu`

DEPARTAMENT DE MATEMÀTICA APLICADA II, UNIVERSITAT POLITÈCNICA DE CATALUNYA, BARCELONA,
SPAIN

E-mail address: `merce.mora@upc.edu`

DEPARTAMENT DE MATEMÀTICA APLICADA III, UNIVERSITAT POLITÈCNICA DE CATALUNYA, BARCELONA,
SPAIN

E-mail address: `ignacio.m.pelayo@upc.edu`

DEPARTAMENT DE MATEMÀTICA APLICADA II, UNIVERSITAT POLITÈCNICA DE CATALUNYA, BARCELONA,
SPAIN

E-mail address: `carlos.seara@upc.edu`

DEPARTAMENT DE MATEMÀTICA APLICADA II, UNIVERSITAT POLITÈCNICA DE CATALUNYA, BARCELONA,
SPAIN

E-mail address: `david.wood@upc.edu`